

SPERNER AND KKM-TYPE THEOREMS ON TREES AND CYCLES

ANDREW NIEDERMAIER, DOUGLAS RIZZOLO, AND FRANCIS EDWARD SU*

ABSTRACT. In this paper we prove a new combinatorial theorem for labellings of trees, and show that it is equivalent to a KKM-type theorem for finite covers of trees and to discrete and continuous fixed point theorems on finite trees. This is in analogy with the equivalence of the classical Sperner's lemma, KKM lemma, and the Brouwer fixed point theorem on simplices. Furthermore, we use these ideas to develop new KKM and fixed point theorems for infinite covers and infinite trees. Finally, we extend the KKM theorem on trees to an entirely new KKM theorem for cycles, and discuss interesting social consequences, including an application in voting theory.

1. INTRODUCTION

Sperner's lemma is a combinatorial theorem about labellings of triangulated simplices that is equivalent to the fixed point theorem of Brouwer and the theorem of Knaster-Kuratowski-Mazurkiewicz (KKM) about covers of simplices by closed sets [7, 9]. In this paper we develop an analogous combinatorial theorem for labellings of trees:

Main Result (see Theorem 1). *Let $T = (V, E)$ be a tree, A a finite subset of V , and let $\ell : V \rightarrow 2^A$ be a proper labelling. Then T contains a fully-labelled edge.*

Our theorem has many useful applications. We show that it is equivalent to a theorem concerning self-maps of trees with no fixed vertices. For metric trees with a finite number of vertices, it also gives an elementary approach to proving a KKM-like theorem and a fixed point theorem (results obtained previously in [3, 6])¹, but our approach has the advantage of being both elementary and, in the case of the fixed point theorem, constructive. Moreover, we show that these theorems are in fact equivalent to our Sperner-type main result. Finally, we use the KKM theorem on metric trees to prove entirely new results for special covers of cycles. Along the way we also consider interesting social interpretations of our results, including an application to voting theory.

2000 *Mathematics Subject Classification.* Primary 05C10; Secondary 55M20, 05C05.

*Corresponding author.

The authors gratefully acknowledge partial support by NSF Grants DMS-0301129, DMS-0701308 (Su) and a Beckman Research Grant at Harvey Mudd College.

¹ The proofs in [6] are for hyperconvex metric spaces, and [1] notes that complete metric trees are hyperconvex metric spaces.

The paper is organized as follows: In Section 2 we prove our Sperner-type theorem about proper labellings of trees. After developing background for metric trees and subdivisions of trees in Section 3, we prove our Tree KKM theorems in Section 4 and give some novel applications in Section 5. We then move to fixed point theorems on finite trees in Section 6, and show there the equivalence of many of our main theorems. We extend our work to infinite trees in Section 7. Finally, we develop a new KKM theorem to cycles in Section 8 and discuss an application to voting theory.

2. A SPERNER-TYPE LEMMA FOR TREES

In this section we prove our main result for labellings of trees. For this result, we view trees as combinatorial (i.e., connected acyclic graphs specified by vertices and edges), although in subsequent sections we shall consider the implications of our result for metric trees (tree-like metric spaces).

Let $T = (V, E)$ be a tree with vertex set V and edge set E . To avoid trivialities, we assume V has at least two vertices. If V is finite, we say T is a *finite* tree; otherwise T is *infinite*. Note that even for an infinite tree, between any two vertices u and w , there is a finite chain of edges that connect u and w and this path of edges is unique.

If v is a vertex, then let $T \setminus v$ denote the graph that results from removing v from V and all the edges incident to v from E . This new graph may have several connected components. Similarly, for an element $e = \{v_1, v_2\} \in E$, we let $T \setminus e$ denote the graph that results from removing e from E . We note that, since T is a tree, $T \setminus e$ has exactly two components.




FIGURE 1. A proper labelling. The non-bracketed numbers mark the vertices that are labels and the bracketed numbers denote the labels of each vertex. The thickened edge is a fully-labelled edge.

Let A be a subset of V which we call the *labels*; every vertex of V will be assigned a collection of labels by a labelling function ℓ . Let 2^A denote the power

set of A , i.e., the set of all subsets of A . A labelling $\ell : V \rightarrow 2^A$ is called a *proper* if:

- for each $a \in A$, $a \in \ell(a)$, and
- for each $v \in V$, the set $A \setminus \ell(v)$, if nonempty, consist of vertices that all lie in one component of $T \setminus v$.

The first condition is actually a consequence of the second, but we stated it separately for clarity. It will be useful to note that the second condition also implies that if e is an edge incident to v , then $\ell(v)$ is missing labels from at most one component of $T \setminus e$. Figure 1 shows an example of a tree with a proper labelling.

Let e be an edge with endpoints x, y . We call that edge *fully-labelled* if $\ell(x) \cup \ell(y) = A$, i.e., if the edge contains all labels. The proper labelling in Figure 1 has a fully-labelled edge. This motivates the statement of our main theorem, which may be viewed as an analogue of Sperner's lemma for trees:

Theorem 1 (Sperner-type Lemma for Trees). *Let $T = (V, E)$ be a tree, let A be a finite subset of V , and let $\ell : V \rightarrow 2^A$ be a proper labelling. Then T contains a fully-labelled edge.*

The finiteness of A is essential, as one may see by considering the *integer tree*: the tree consisting of a line whose vertices are the integers, and whose edges are intervals between successive integers (see Figure 3). Now let $A = V$ and label each integer n by all the integers less than or equal to n . This labelling is proper, but it does not have a fully-labelled edge.

Proof. It suffices to consider the case where V is also finite, for if not, we may restrict our attention to the subtree of T spanned by A —this is a finite tree because A is finite, and any fully-labelled edge in this subtree is fully-labelled in T . So assume V is finite.

If there exists v in V with $\ell(v) = A$, then desired conclusion holds: any edge containing v is fully-labelled. Otherwise, suppose there is no such vertex. We shall construct a “successor” function $S : V \rightarrow V$ which will be useful for locating a fully-labelled edge. For each v in V , since $\ell(v) \neq A$, $\ell(v)$ is missing labels from exactly one component of $T \setminus v$. In that component, let $S(v)$ be the unique vertex that shares an edge with v in T .

Let α be a leaf of T . Consider the sequence $\{v_n\}$ given by $v_1 = \alpha$ and $v_k = S(v_{k-1})$ for $k > 1$. Since V is finite, this sequence must be eventually periodic. Since T is a tree, this period cannot be of length greater than two. However, if $S(x) = y$ and $S(y) = x$, then x, y are endpoints of an edge e that we claim is fully-labelled. Why?

The vertices missing from $\ell(y)$ are in the component of $T \setminus y$ containing x (since $S(y) = x$), and the vertices missing from $\ell(x)$ are in the component of $T \setminus x$ containing y (since $S(x) = y$). Then the vertices missing from $\ell(x)$ and the vertices missing from $\ell(y)$ are from different components of $T \setminus e$. So there are

no vertices of A that are missing from both $\ell(x)$ and $\ell(y)$, i.e., $\ell(x) \cup \ell(y) = A$ and e is a fully-labelled edge. $\square$

Note that the above successor function gives rise to a method for locating a fully-labelled segment, using the sequence of vertices obtained by iterating S until one reaches a vertex for which S is not defined (and therefore has all labels), or until the sequence repeats.

Theorem 1 gives rise to an interesting corollary for functions on vertices of a tree:

Theorem 2. *Let $T = (V, E)$ be a finite tree and $f : V \rightarrow V$ a function. Then either f fixes some vertex, or there is an edge $e \in E$ with endpoints x and y such that e is on the path from $f(x)$ to $f(y)$.*

Proof. Suppose f does not fix any vertex. Consider a labelling $\ell : V \rightarrow 2^V$ such that for each vertex v , $\ell(v)$ is the set containing v together with the vertices of T that are not in the same component of $T \setminus v$ as $f(v)$. This is well-defined and a proper labelling because f does not fix any vertices. Theorem 1 now implies that there is a fully-labelled edge e with endpoints x and y . Removing e from T produces a graph with exactly two non-empty components.

If $f(x)$ and $f(y)$ are in the same component of $T \setminus e$, then the path from $f(x)$ to $f(y)$ does not contain e . So at least one of x and y is not on the path from $f(x)$ to $f(y)$; suppose without loss of generality it is x . Then $f(x)$ and $f(y)$ are in the same component of $T \setminus x$, so by the definition of ℓ , $\ell(x)$ does not contain $f(y)$. Since $\ell(y)$ clearly does not contain $f(y)$ either, this contradicts the fact that e is fully-labelled.

Therefore $f(x)$ and $f(y)$ are in different components of $T \setminus e$, which implies that e is on the unique path from $f(x)$ to $f(y)$. $\square$

Theorem 2 may be viewed as a kind of “discrete” fixed point theorem for trees, because it says that if f fixes no vertices, then some edge must intersect the span of the images of its endpoints. Compare it to the continuous version we prove later in Theorem 8. Moreover,

Theorem 3. *Theorem 2 is equivalent to Theorem 1.*

Proof. Having already shown Theorem 1 implies Theorem 2, we now show the converse. Suppose that $\ell : V \rightarrow 2^A$ is a proper labelling of T . If there is a vertex v such that $\ell(v) = A$, then any edge containing v is fully-labelled. Otherwise, for each vertex v define $f(v)$ to be any label which is not in $\ell(v)$. Clearly f is a function with no fixed points, so Theorem 2 implies that there is an edge $e = \{x, y\}$ such that e is on the path from $f(x)$ to $f(y)$.

We claim that e is fully-labelled. Since $f(x)$ and $f(y)$ are in different components of $T \setminus e$, then the definition of f shows that $\ell(x)$ and $\ell(y)$ are missing labels from different components of $T \setminus e$. So there are no vertices of A that are missing from both $\ell(x)$ and $\ell(y)$, i.e., $\ell(x) \cup \ell(y) = A$ and e is a fully-labelled edge. $\square$

3. METRIC TREES AND SEGMENTATIONS

Theorems 1 and 2 have several applications to metric trees, which are essentially combinatorial trees that are realized as metric spaces by replacing edges by line segments isometric to a compact interval of $\mathbb{R}$. For the sake of clarity, we make precise in this section what we mean by *metric tree* as well as the concept of a subdivision called a *segmentation*, but there are no surprises here, so this brief section may be skimmed if desired.

A *metric tree* may be viewed as a triple $T = (V, E, X)$, with a vertex set V (that may be finite or infinite), an edge set E , and *underlying metric space* X . Here, (V, E) specifies a combinatorial tree and the metric space X has the following relationship with V and E . For each edge $e \in E$, let the *realized edge* e_X be an isometric copy of a segment of the real line $[0, L_e]$ (via some isometry $\phi_e : [0, L_e] \rightarrow e_X$); then e has a metric and L_e is the length of the edge e under this metric.

Let X be the quotient set obtained by identifying endpoints of realized edges with the vertex set V according to the instructions in E ; e.g., if an edge $e \in E$ is the pair $\{v, w\}$, then $\phi_e(0)$ in e_X is identified one element of e and $\phi_e(L_e)$ in e_X is identified with the other. In this way, realized edges are “glued” together at endpoints just as they are in the combinatorial tree (V, E) .

Since (V, E) has no cycles, between any two points x, y in the metric space X , there is a unique (non-self-intersecting) path between x and y . X has a natural metric: $d(x, y)$ defined to be the length of the path between x and y , i.e., the sum of the lengths of the edges (or partial edges) along this unique path. It will be useful to note if z is on the path between x and y , then the triangle inequality becomes an equality: $d(x, y) = d(x, z) + d(z, y)$.

We remark that, unlike other definitions of metric trees in the literature, our definition of metric tree retains the combinatorial structure specified by V and E .

Given a tree $T = (V, E, X)$ it will be convenient at times to consider a *segmentation* of T , which is another metric tree obtained from T by finite subdivision of its edges. In particular, $T' = (V', E', X)$ is a *segmentation* of $T = (V, E, X)$ if: (1) $V' = V \cup V^*$ where V^* is a collection of points $\{v_\alpha\}_{\alpha \in A}$ from X so that at most a finite number of the v_α come from a realized edge e_X , $e \in E$, and (2) E' is the collection of edges obtained from E in the natural way (by deleting edges in which elements of V^* appear and including edges of the implied subdivision along that edge). Note that the metric spaces for T' and T are the same, so the set of continuous functions on T and T' are the same.

Recall that the diameter of a set S is given by $\text{diam}(S) = \sup_{x, y \in S} d(x, y)$. With this, define the *size* of a segmentation $T' = (V', E', X)$ by $\text{size}(T') = \max_{e \in E'} \text{diam}(e_X)$. Clearly every tree has an arbitrarily small segmentation. Note also that every point in a metric tree T that is not a leaf is a *cut point*: its removal “cuts” T into more than one path-connected component.

In what follows, all trees T are metric trees.




FIGURE 2. A KKM cover of a tree relative to the vertices $\{1, 2, 4, 6\}$. The sets of a KKM cover do not have to be connected sets (although they are in this diagram).

4. KKM COVERS OF TREES

Just as the traditional Sperner's Lemma implies the classical KKM theorem for covers of simplices by closed sets, Theorem 1 implies a KKM theorem for metric trees. While this KKM theorem can be derived as a corollary of the results in [3] and [6], our proof is more elementary.

Let $T = (V, E, X)$ be a tree, and A a subset of points in X . We shall call a family of sets $\mathcal{D} = \{D_a : a \in A\}$ a *KKM cover of T relative to A* if:

- each D_a is a closed subset of X ,
- each $a \in D_a$, and
- for any two points a and b in A , the path between them is contained in $D_a \cup D_b$.

We call the last condition the *path-covering property* of $\mathcal{D}$; it is analogous to the facet-covering property of KKM covers of simplices. So each point in A has an associated closed set in $\mathcal{D}$ that contains it, and the path between any two points in A is covered by the corresponding sets. See Figure 2.

If $A = V$, we may simply say $\mathcal{D}$ is a *KKM cover* of T , and the sets of $\mathcal{D}$ will cover the underlying metric space X . (Note that if $A \neq V$, then a KKM cover of T relative to A may not cover X , but it will cover the subtree spanned by A .)

The following lemma defines a labelling that is associated to each KKM cover relative to a subset of *vertices*:

Lemma 4. *Let $T = (V, E, X)$ be a tree, let $A \subset V$ be a subset of vertices, and let $\mathcal{D} = \{D_a : a \in A\}$ be a KKM cover of T relative to A . The labelling $\ell : V \rightarrow 2^A$ defined by $\ell(v) = \{a : v \in D_a\}$ is a proper labelling.*

We call this labelling ℓ the *membership labelling* associated with $\mathcal{D}$, because the labelling identifies all the sets in $\mathcal{D}$ that contain v and assigns the corresponding vertices of A as labels. (Note that if v is not covered by the sets of $\mathcal{D}$, then $\ell(v)$ may be empty.) Note that A may be finite or infinite.

As an example, in Figure 2, all vertices in the set D_1 will have 1 in their label set. Similarly, because vertex 4 is in sets D_2 and D_4 and D_6 , $\ell(4)$ will contain 1, 2, and 4. The leaf at top left will have an empty label set. The reader may notice that the labelling in Figure 1 is the membership labelling for the KKM cover in Figure 2 if labels 3 and 5 were removed from every label set.

Proof. To show that $a \in \ell(a)$ for all $a \in A$, note that $a \in D_a$ because $\mathcal{D}$ is a KKM cover relative to A .

To show the second condition for a proper labelling, fix a vertex v in V , and suppose there were two vertices a and b in A that are not in $\ell(v)$. Then both sets D_a and D_b do not contain v . If a and b were in different components of $T \setminus \{v\}$, then v must lie on the unique path between a and b in T and the path-covering property of $\mathcal{D}$ would imply $v \in D_a \cup D_b$, a contradiction. Therefore any vertices not in $\ell(v)$ must lie in one component of $T \setminus \{v\}$, as desired. $\square$

We are now prepared to prove a KKM-like theorem for finite covers of trees:

Theorem 5 (Tree KKM Theorem). *Let $T = (V, E, X)$ be a metric tree, A a finite subset of points of X , and suppose $\mathcal{D} = \{D_a : a \in A\}$ is a KKM cover of T relative to A . Then*

$$\bigcap_{a \in A} D_a \neq \emptyset.$$

Proof. We may as well assume that V is finite, for otherwise we may restrict our attention to the subtree K spanned by a finite set of edges that contain A , which contains a finite number of vertices. Any KKM cover of T relative to A will also restrict to a KKM cover of K relative to A , and a nonempty intersection of the KKM cover of K would imply a nonempty intersection of the KKM cover of T .

Suppose, by way of contradiction, that the intersection $\bigcap_{a \in A} D_a$ were empty. Then the set of complements $\mathcal{C} = \{T \setminus D_a : a \in A\}$ is an open cover of T . Since V is finite, X is compact and this cover has a Lebesgue number δ . Let $T' = (V', E', X)$ be a segmentation of T with $\text{size}(T') < \delta$ chosen so that A is a subset of the vertices of V' .

Consider the membership labelling $\ell : V' \rightarrow 2^A$ defined by $\ell(v') = \{a : v' \in D_a\}$. Lemma 4 shows that ℓ is a proper labelling. By Theorem 1, there exists an edge $e \in T'$ with endpoints x and y such that $\ell(x) \cup \ell(y) = A$. Thus, for all $a \in A$, either $x \in D_a$ or $y \in D_a$ (or both). However, since $\text{size}(T') < \delta$, the Lebesgue number property guarantees that $e \subseteq T \setminus D_a$ for some a , implying that $e \cap D_a = \emptyset$, a contradiction. Therefore, we conclude that $\bigcap_{a \in A} D_a \neq \emptyset$. $\square$

Note also that the sets of a KKM cover do not have to be connected (though they are in Figure 2. However, if a tree is covered by sets that are connected as well as pairwise intersecting, then it is a KKM cover!

Corollary 6 (Tree KKM for Connected Sets). *Let $\mathcal{D} = \{D_1, D_2, \dots, D_k\}$ be a finite collection of closed, connected sets that cover a metric tree $T = (V, E, X)$ such that each pair $D_i \cap D_j$ is nonempty. Then there is a point x in all the sets of $\mathcal{D}$.*

Proof. Choose points $a_i \in D_i$ for each i , and put them in a set A . To show $\mathcal{D}$ is a KKM cover of T relative to A , it remains to show the path-covering property.

If for some pair a_i and a_j in A , the path-covering property did not hold, then the path between a_i and a_j would contain a point y that is not covered by $D_i \cup D_j$. Then $X \setminus \{y\}$ would have two connected components that would separate a_i from a_j , so D_i and D_j would then lie in different components and could not be pairwise intersecting, a contradiction. $\square$

5. APPLICATIONS

In this section we indicate some applications of Theorem 5. We remark that our proof of Theorem 1 suggests associated constructive algorithms for finding the locations suggested by both of these examples.

Pizza Delivery. Suppose you are starting a pizza delivery business and you desire a good location for your store. Your city has several neighborhoods connected by a tree of roads. For each neighborhood i , there is a “deliverability” set D_i : the set of all locations on the tree with an acceptable commute to neighborhood i . These sets are naturally closed and connected, as in Figure 2. Then the Tree KKM Theorem for Connected Sets (Corollary 6) says that if for every pair of neighborhoods i and j there is a common acceptable location to place your store, then there will be a location with an acceptable commute to all neighborhoods.

Grand Central Station. Suppose several cities are connected by a tree of train tracks. Each city has its own train authority, and suppose it is possible to get from city i to city j using only those cities’ trains (possibly switching several times). Then Theorem 5 shows that there must be a location through which trains from all cities must pass, i.e., there is a location where one could place a Grand Central Station.

Note that something further is true if we make some intuitive assumptions about the structure of the tree of train tracks. It seems reasonable to suppose that each vertex of this tree is a station and that trains only change directions at stations. With these assumptions the following result becomes apparent: there must already be a station at which trains from all of the cities stop, i.e., a Grand Central Station already exists. If the point guaranteed by Theorem 5 is not a station, then trains from each city also must pass through the two nearest stations on either side of this point (because trains only change directions at stations).

6. A FIXED POINT THEOREM FOR FINITE TREES

Just as the KKM Theorem and Sperner’s Lemma imply Brouwer’s Theorem on simplices, we can use our previous theorems to prove a classical fixed point

theorem for trees. We will give two proofs that every continuous function from a finite tree to itself has a fixed point. The first proof will use our KKM Theorem and the second will use Theorem 1. Both proofs are provided because they have different strengths. The KKM approach is short and sweet, providing a quick, clean proof of the theorem while the approach of Theorem 1 gives an algorithm for locating a fixed point.

We shall often appeal to a special family of closed sets associated to any continuous self-map on a tree. Given $T = (V, E, X)$ be a metric tree, and $A \subset V$ a subset of vertices, and $f : T \rightarrow T$ be a continuous function, let $\mathcal{D}_{f,A} = \{D_a : a \in A\}$ be the family of sets defined by

$$(1) \quad D_a = \{x \in T : d(x, a) \leq d(f(x), a)\}.$$

To interpret, D_a contains the set of all points that do not move closer to a after applying the function f , i.e., all points that stay fixed or move away from a . As we show next, this family $\mathcal{D}_{f,A}$ is a KKM cover, so we shall sometimes call it the *move-away KKM cover associated to f and A* .

Theorem 7. *The family $\mathcal{D}_{f,A}$ defined above is a KKM cover of T relative to A .*

Proof. It should be apparent from the definition that $a \in D_a$ for each $a \in A$.

We now show that each D_a is closed. Let $\{x^k\}$ be a sequence in D_a that converges to x in T . Since f is continuous, we see that $f(x^k) \rightarrow f(x)$. Let $\epsilon > 0$ be given. Choose N such that $k \geq N$ implies that $d(x^k, x) < \epsilon$ and $d(f(x^k), f(x)) < \epsilon$. Then

$$\begin{aligned} d(x, a) &\leq d(x, x^k) + d(x^k, a) \\ &\leq d(x, x^k) + d(f(x^k), a) \\ &\leq d(x, x^k) + d(f(x^k), f(x)) + d(f(x), a) \\ &< 2\epsilon + d(f(x), a). \end{aligned}$$

Since this is true for all $\epsilon > 0$ we conclude that $d(x, a) \leq d(f(x), a)$, so $x \in D_a$. Therefore D_a is closed.

Finally, we show the path-covering property of $\mathcal{D}_{f,A}$: if a and b are in A , and if x is on the path between them, we shall show that $x \in D_a \cup D_b$. Suppose that $f(x) = x$, then clearly $x \in D_a$ and $x \in D_b$ as desired. Otherwise, suppose that $f(x)$ is not in the same component of $T \setminus \{x\}$ as a . Then since T is a tree there is only one path from a to $f(x)$ and it must contain x . It follows that the path from a to x is contained in the path from a to $f(x)$, so we conclude that $d(a, x) \leq d(a, f(x))$, hence $x \in D_a$. By similar argument, if $f(x)$ is not in the same component of $T \setminus \{x\}$ as b , then $x \in D_b$. Either way, $x \in D_a \cup D_b$. $\square$

We can now give our first proof of the fixed point theorem for trees with a finite number of vertices:

Theorem 8. *(The Tree Fixed Point Theorem) Let $T = (V, E, X)$ be a metric tree, V be finite, and $f : T \rightarrow T$ be a continuous function. Then f has a fixed point.*

First Proof of Theorem 8. Consider the family $\mathcal{D}_{f,V}$ defined in (1). Theorem 7 shows that it is a KKM cover of T . Since V is finite, Theorem 5 shows that there is some point x in the intersection of all sets of $\mathcal{D}_{f,V}$. By definition of $\mathcal{D}_{f,V}$, we have that for all $v \in V$, $d(x, v) \leq d(f(x), v)$.

Suppose that $x \neq f(x)$. If $f(x)$ were a leaf, say v , this would contradict that $x \in D_v$. So $f(x)$ is not a leaf and is thus a cut point. Choose $w \in V$ such that w and x are in different components of $T \setminus \{f(x)\}$. Then the path from w to $f(x)$ is contained in the path from w to x . Since $f(x) \neq x$, this implies that $d(x, w) > d(f(x), w)$, so that $x \notin D_w$, a contradiction. Hence $f(x) = x$. $\square$

Before giving the second proof of the theorem, we need a couple of additional lemmas.

If a, b are two points in $T = (V, E, X)$, let $[a, b]$ denote the set of all points on the unique path between a and b in X , and let d denote the metric on X .

Lemma 9. *Let $T = (V, E, X)$ be a tree and suppose that $f : T \rightarrow T$ is continuous. Suppose that $T' = (V', E', X)$ is a segmentation of T . Let $\ell : V' \rightarrow 2^V$ be defined by*

$$\ell(v') = \{v \in V : d(v', v) \leq d(f(v'), v)\}.$$

If e is a fully-labelled edge with endpoints y and z , then either $[y, z] \cap f([y, z]) \neq \emptyset$, or $d(y, f(z)) \leq d(f(y), f(z))$.

Said another way, the labelling $\ell(v')$ specifies the vertices v in V for which the direction of motion from v' to $f(v')$ is fixed or away from v .

Proof. Suppose that $[y, z] \cap f([y, z]) = \emptyset$. Since f is continuous and $[y, z]$ is path-connected, $f([y, z])$ is also path-connected and thus contained in a single component of $T \setminus [y, z]$. Let $v \in V$ be a leaf in a component of $T \setminus \{f(z)\}$ that does not contain $[y, z]$ (if $f(z)$ is itself a leaf let $v = f(z)$). Then since $f(z)$ separates v from z , $f(z)$ must be on the path from v to z . So $f(z)$ is closer to z than v is to z , hence $v \notin \ell(z)$. So $v \in \ell(y)$ because e is fully-labelled. This means that $f(y)$ is not closer to v than y is to v , i.e., $d(v, y) \leq d(v, f(y))$.

But $f(z)$ is on the path from v to y , since it separates v from y . Thus:

$$d(v, f(z)) + d(f(z), y) = d(v, y) \leq d(v, f(y)) \leq d(v, f(z)) + d(f(z), f(y)),$$

which implies that $d(y, f(z)) \leq d(f(y), f(z))$, as desired. $\square$

The following lemma is standard but we include it for completeness.

Lemma 10 (Epsilon Fixed Point Theorem). *Suppose that K is a compact subset of the metric space (X, d) and that $f : K \rightarrow K$ is continuous. If f has an ϵ -fixed point for every $\epsilon > 0$ then f has a fixed point.*

Proof. Let $\{a_n\}$ be a sequence of $1/n$ -fixed points, that is, $d(a_n, f(a_n)) < 1/n$ for all n . Since K is compact it is sequentially compact and thus $\{a_n\}$ has a convergent subsequence, which we denote $\{a'_n\}$ with $a'_n \rightarrow x \in K$. Let $\epsilon > 0$.

Since $a'_n \rightarrow x$ there exists N_1 such that $n \geq N_1$ implies that $d(a'_n, x) < \epsilon/2$. Let $N = \max(N_1, 2/\epsilon)$. Then $n \geq N$ implies that

$$d(x, f(a'_n)) \leq d(x, a'_n) + d(a'_n, f(a'_n)) < \epsilon,$$

so that $f(a'_n) \rightarrow x$. However, since f is continuous, we have also that $f(a'_n) \rightarrow f(x)$. Hence $f(x) = x$, and x is a desired fixed point. $\square$

With these lemmas in hand, we are prepared to give the second proof of the fixed point theorem, which shows how to find approximate fixed points.

Second Proof of Theorem 8. By Lemma 10, it is sufficient to show that T has an ϵ -fixed point for all $\epsilon > 0$. Let $\epsilon > 0$ be given. Since T is compact, f is uniformly continuous, so we let $\delta_1 > 0$ be such that if x and y are points in T with $d(x, y) < \delta_1$ then $d(f(x), f(y)) < \epsilon/2$. Let $\delta = \min(\delta_1, \epsilon/2)$ and let $T' = (V', E', X)$ be a segmentation of $T = (V, E, X)$ with $\text{size}(T') < \delta$.

As above, let $\ell : V' \rightarrow 2^V$ be defined by $\ell(v') = \{v \in V : d(v', v) \leq d(f(v'), v)\}$. Note that this is just the membership labelling defined in Lemma 4 for the move-away KKM cover $\mathcal{D}_{f,A}$ defined at the start of this section. Thus by Theorem 1 there is a fully-labelled segment with endpoints y and z .

Let w be a point in $f([y, z]) \cap [y, z]$ if this intersection is non-empty or let $w = f(z)$ if the intersection is empty. We see that $d(y, f(y)) \leq d(y, w) + d(w, f(y))$. If $w \in f([y, z]) \cap [y, z]$ then $d(y, w) < \delta$ and $d(w, f(y)) < \epsilon/2$, and if $w = f(z)$ then by Lemma 9 we have that $d(y, f(z)) \leq d(f(y), f(z)) < \epsilon/2$. In either case we have that

$$d(y, f(y)) \leq d(y, w) + d(w, f(y)) < \frac{\epsilon}{2} + \frac{\epsilon}{2}.$$

Therefore y is an ϵ -fixed point. $\square$

The constructive nature of Theorem 1 and this proof suggests a method for locating an approximate fixed point. Namely, choosing a segmentation with a sufficiently small size and using the labelling ℓ defined above, start at a leaf and “move in the direction of the missing labels” as suggested by the successor function S in the proof of Theorem 1. This will eventually lead to a fully-labelled edge whose endpoints are approximate fixed points.

We note that Theorem 8 implies the Tree Discrete Fixed Point Theorem (Theorem 2), and thus:

Theorem 11. *The following are all equivalent:*

- (1) *Sperner-type Lemma for Trees (Theorem 1),*
- (2) *Tree Discrete Fixed Point Theorem (Theorem 2),*
- (3) *Tree KKM Theorem (Theorem 5), and*
- (4) *Tree Fixed Point Theorem (Theorem 8).*

Proof. Theorem 3 showed that (2) and (1) are equivalent. We have also seen that (1) implies (3), and (3) implies (4), so it suffices to show that (4) implies (2).

Let (V, E) be a finite tree and $f : V \rightarrow V$ a function. Construct a metric tree $T = (V, E, X)$ from (V, E) in the natural way, by realizing E as edges of length

1. We now construct a continuous function $\hat{f} : X \rightarrow X$ by extending the given f linearly across single edges, i.e., if $\{v, w\}$ is an edge in E and x is a point on the realized edge that is a fraction of distance r along the path from v to w , then let $\hat{f}(x)$ be the point that is a fraction of distance r along the path from $f(v)$ to $f(w)$. By construction, this function has the property that $f(v)$ and $f(w)$ must be in different components of $X \setminus \{x\}$.

By Theorem 8, there is a point z such that $\hat{f}(z) = z$. If $z \in V$ then we see f has a fixed vertex, as desired. Otherwise, z is on some edge $e = \{v, w\}$ and as noted above, $f(v)$ and $f(w)$ must be in different components of $X \setminus \{z\}$; moreover, they must be vertices in V . But v and w are the nearest vertices to z in those two components. Hence the path from $f(v)$ to $f(w)$ is a path between vertices of V that contains e . $\square$

The equivalence of our results for trees is in analogy with the classical equivalence of the classical Sperner's Lemma and KKM theorem and the Brouwer fixed point theorem.

7. INFINITE TREES

With additional hypotheses, we can extend both the Tree KKM theorem and the Tree Fixed Point Theorem to a tree T to the infinite setting (for covers with infinitely many sets in Theorem 5, and to trees with infinitely many vertices in Theorem 8). When T is compact, our methods produce results previously shown in [6]. However, we allow T to be non-compact as well, as long as the map f is compact.

The following lemma will be useful:

Lemma 12. *Let X be a topological space. Let Λ be an infinite (not necessarily countable) index set and suppose that $\mathcal{D} = \{D_\alpha : \alpha \in \Lambda\}$ is a family of closed sets in X with the finite intersection property. Further suppose that D_α is compact for at least one $\alpha \in \Lambda$. Then*

$$\bigcap_{\alpha \in \Lambda} D_\alpha \neq \emptyset.$$

Proof. If not, then there is no point in every D_α , hence each point is covered by some complement: D_α^c . Then $\mathcal{D}^c = \{D_\alpha^c : D_\alpha \in \mathcal{D}\}$ is an open cover of $Y = \bigcup \{D_\alpha : \alpha \in \Lambda\}$. Let D_β be one of the compact sets in $\mathcal{D}$. It follows that $\mathcal{D}^c$ is an open cover of D_β and thus has a finite subcover, say it is the collection $\mathcal{F}$. Since these sets cover D_β , the intersection of their complements together with D_β is empty — that is,

$$\left(\bigcap_{D_\alpha^c \in \mathcal{F}} D_\alpha \right) \cap D_\beta = \emptyset.$$

This contradicts that $\mathcal{D}$ has the finite intersection property and thus proves the lemma. $\square$

Now we may prove a KKM theorem for infinite trees:

Theorem 13 (KKM Theorem for Infinite Trees). *Let $T = (V, E, X)$ be a tree, let A be a (possibly infinite) subset of V . Suppose that $\mathcal{D} = \{D_a : a \in A\}$ is a KKM cover of T relative to A such that at least one set in $\mathcal{D}$ is compact. Then*

$$\bigcap_{a \in A} D_a \neq \emptyset.$$

The extra condition that one set be compact is essential. Recall again the integer tree: a line whose nodes are the integers, and whose edges are the intervals between successive integers. See Figure 3. One may construct a KKM cover by letting $D_i = [i, \infty)$, but the intersection of all such sets is empty.




FIGURE 3. The integer tree, with infinite number of vertices (at integers) and edges.

Proof. By Lemma 12 it suffices to show that $\mathcal{D}$ has the finite intersection property. Let J be a finite subset of A . Then the family of sets $\{D_j : j \in J\}$ forms a KKM cover of T relative to J , a finite set, so Theorem 5 applies. Hence $\bigcap_{\alpha \in J} D_\alpha \neq \emptyset$. $\square$

Before proving our fixed point theorem for infinite trees, we need the following lemmas:

Lemma 14. *If T is a tree, and $C \subseteq T$ is connected, then C is path-connected.*

Proof. Suppose $x, y \in C$ but are not connected by a path in C . They are, however, connected by a path in T , because T is a tree. Then there is a point p on the path that is not in C . Then $T \setminus \{p\}$ has at least two components, one of which contains x and another of which contains y . Furthermore, each of these components is open as a subset of T . Let A be the component containing x and B be the union of all other components. Then $A \cap C$ and $B \cap C$ provide a disconnection of C . $\square$

This theorem yields an easy corollary:

Corollary 15. *If T is a tree and $C \subseteq T$ is path-connected then $\overline{C}$ is path-connected.*

Proof. If C is path-connected, then it is connected. Then $\overline{C}$ is connected, and by the previous proposition, $\overline{C}$ is path-connected. $\square$

Using Theorem 5 we can prove the following fixed point theorem. Recall that if X and Y are metric spaces, $f : X \rightarrow Y$ is a *compact* map if $f(B)$ is contained in a compact subset of Y for every bounded set B .

Theorem 16. *Suppose that T is a bounded tree with vertex set V . If $f : T \rightarrow T$ is a continuous compact map, then f has a fixed point.*




FIGURE 4. A bounded infinite tree. Vertices (not all shown) accumulate at the open endpoints of the underlying set.

We give a couple of (non-)examples to illustrate. Consider the integer tree of Figure 3 with the map that shifts it one unit to the right. This map does not have a fixed point; it is a compact map, but the tree is not bounded. The infinite tree of Figure 4 is bounded, but it is not compact. One may easily construct a fixed-point-free map that moves points within this tree towards the rightmost open endpoint.

Proof. Since f is compact and continuous, by Corollary 15, $K = \overline{f(T)}$ is compact and connected. Let ∂K denote the boundary of K in T . Consider $V^* = (V \cap K) \cup \partial K$, the set of all vertices of T that are in K , as well as the boundary points of K in T . It is a subset of K since K is compact.

We shall construct a new tree T^* with vertices V^* and underlying metric space K . To start, let T° be the subtree of T spanned by the vertices in $V' = V \cap K$, with edge set E' . We shall augment T° to form T^* by adding points of ∂K as leaves.

So consider any $x \in \partial K$. Note that x lies in some realized edge e of T with endpoints in V .

We claim that exactly one component C of $e \setminus \{x\}$ will intersect $f(T)$. This follows because x is an endpoint of e , or otherwise from the fact that $f(T)$ is connected (if $s, t \in f(T)$ but s, t were in different components, then the path from s to t would be completely in $f(T)$ and x could not be a boundary point of K).

This component C contains an endpoint, call it v_x . Then either v_x is in $f(T)$ or not. We explore these cases and the tree T^* that results:

- (1) If $v_x \notin f(T)$, then because $f(T)$ is connected, then $f(T)$ must be completely contained within the interior of e , it is an interval with x at one

endpoint (because $x \in K = \overline{f(T)}$) and $y \neq v_x$ at another endpoint. Then K has at most two boundary points (one if $x = v_x$) and is an interval. Then let T^* be the tree consisting of one edge $e^* = [x, v_x]$ and two vertices.

- (2) If $v \in f(T)$, then $f(T)$ contains C because $x \in K = \overline{f(T)}$. Then the segment $e_x^* = [x, v_x]$ lies completely in K , and $v_x \in V' = V \cap K$. This construction may be done for every $x \in \partial K$. We construct T^* as the tree with vertex set $V^* = V' \cup \{v_x : x \in \partial K \setminus V\}$, edge set $E^* = E' \cup \{e_x^* : x \in \partial K \setminus V\}$, and underlying space K .

Since T^* has underlying space K , note that $f : T \rightarrow T$ restricts to a function $f^* : T^* \rightarrow T^*$. Then consider the move-away KKM cover $\mathcal{D}_{f^*, V^*}$ defined in (1); it is a KKM cover of T^* relative to V^* .

Suppose that H is a finite subset of V^* . From Theorem 7 we see that $\mathcal{D}_{f^*, H}$ is a KKM cover of T relative to H and thus, by Theorem 5, the intersection of its sets is nonempty. Thus, $\mathcal{D}_{f^*, A^*}$ has the finite intersection property.

Because K is compact, all the sets of $\mathcal{D}_{f^*, A^*}$ are compact, so Theorem 13 shows that the intersection of sets in $\mathcal{D}_{f, A}$ is also non-empty, say it contains a point z . Then for all $v \in V^*$ we have $d(z, v) \leq d(f(z), v)$. We claim that $f(z) = z$.

Suppose that $z \neq f(z)$. If $f(z)$ were a leaf, say v , this would contradict that $z \in D_v$. So $f(z)$ is not a leaf and is thus a cut point. Choose $w \in V$ such that w and z are in different components of $T \setminus \{f(z)\}$. Then the path from w to $f(z)$ is contained in the path from w to z . Since $f(z) \neq z$, this implies that $d(z, w) > d(f(z), w)$, so that $z \notin D_w$, a contradiction. Hence $f(x) = x$. $\square$

The compactness hypothesis is reminiscent of the compactness hypothesis of the Schauder fixed point theorem [4].

8. A KKM THEOREM FOR CYCLES

Recall that a *cycle* is a finite graph with vertices $v_1, \dots, v_n$ and edges (v_i, v_{i+1}) as well as (v_n, v_1) . We define a *metric cycle* to be a triple $C = (V, E, X)$ where (V, E) is a cycle (as above) and X is an *underlying metric space* obtained from a cycle in exactly the same fashion as we obtained a metric tree from a tree in Section 3. The resulting space C is topologically a circle, partitioned into a finite set of segments (realized edges) joined at their endpoints $v_1, \dots, v_n$. Between any two points of C there are exactly two paths; the metric is just the minimum length of the two paths connecting the points.

Hereafter, all metric cycles will simply be referred to as cycles.

If C_n is a cycle with n vertices and x is in C_n , let $e(x)$ be the set consisting of x and all points y that are not vertices but are on a realized edge with x . Note that by removing from C_n the set $e(x)$ as well as vertices and realized edges within, we obtain a metric tree $C_n \setminus e(x)$ with n or $(n-1)$ vertices, depending on whether or not x is a vertex of C_n . This observation will become the key to reducing KKM covers on cycles to KKM covers on trees.

Now, since there are two paths connecting any two distinct vertices in a cycle, we must slightly alter our definition of KKM cover for trees, but we want to do so in a way consistent with our definition for trees.

With this in mind, let C_n be a cycle with n vertices $V = \{v_1, v_2, \dots, v_n\}$. A *KKM cover* of C_n is a family of sets $\mathcal{D} = \{D_v : v \in V\}$ such that the following conditions hold:

- each D_v is closed,
- each $v \in D_v$, and
- for all $v, w \in V$, at least one of the paths between v and w is contained in $D_v \cup D_w$.

This new path-covering property generalizes the corresponding property for trees. We can now state the main theorem of this section.

Theorem 17. *Suppose that C_n is a cycle with vertices $V = \{v_1, v_2, \dots, v_n\}$ and let $\mathcal{D} = \{D_v : v \in V\}$ be a KKM cover of C_n . Then there is a point x in C_n such that x is in at least $\lfloor \frac{n}{2} \rfloor + 1$ sets of $\mathcal{D}$.*

To compare this result with Theorem 5, note that KKM covers of trees have a point in *all* the sets of the cover, but KKM covers of cycles have a point in a *strict majority* of the sets.

Proof. For each x one may consider the set $\ell(x) = \{v \in V : x \in D_v\}$. If there were an x such that the cardinality $|\ell(x)| \geq \lfloor \frac{n}{2} \rfloor + 1$, we would be done, so assume that for every x , either $|\ell(x)| = \lfloor \frac{n}{2} \rfloor$ or $|\ell(x)| \leq \lfloor \frac{n}{2} \rfloor - 1$.

If $|\ell(x)| \leq \lfloor \frac{n}{2} \rfloor - 1$, then let $H = V \setminus \ell(x)$; clearly $|H| \geq \lfloor \frac{n}{2} \rfloor + 1$. Note that for any pair v and w in H , the path between them covered by $D_v \cup D_w$ does not include x , so this path is still covered if we remove $e(x)$ from the graph. Then $C_n \setminus e(x)$ is a tree and the family $\mathcal{F} = \{D_v : v \in H\}$ is a KKM cover of this tree relative to H . By Theorem 5, $\mathcal{F}$ has non-empty intersection, and it has at least $\lfloor \frac{n}{2} \rfloor + 1$ sets of $\mathcal{D}$, as desired.

Finally, consider the case in which $|\ell(x)| = \lfloor \frac{n}{2} \rfloor$ is constant for all x . We now show why this leads to a contradiction.

The set of boundary points of D_v , denoted by ∂D_v , is a closed set. Moreover, since D_v is closed, ∂D_v has no interior (since any neighborhood of $x \in \partial D_v$ must contain a point $p \notin D_v$, hence also $p \notin \partial D_v$). But then the union of boundary points $B = \cup \{\partial D_v : v \in V\}$ has no interior, because it is the finite union of closed sets with no interior (this can be proved by induction). Also, B is closed because all the ∂D_v are. So B^c is a nonempty open set.

So choose $x \in B^c$ and since B is closed, we may find $b \in B$ that is closest to x . Let $U = \{y : d(x, y) < d(x, b)\}$, so all points in U are, for each $v \in V$, interior points of either D_v or D_v^c . So if $y \in U$, $\ell(x) = \ell(y)$. Since b is a limit point of U and each D_v is closed, $\ell(x) \subseteq \ell(b)$.

If for some $v \notin \ell(x)$ we have $b \in D_v$, then $\ell(x) \cup \{v\} \subseteq \ell(b)$, so the size of ℓ is not constant, a contradiction. Otherwise, for every $v \notin \ell(x)$, we have that $b \in D_v^c$, an open set, so $\ell(x) = \ell(b)$. So there is an open set W around b that

contains no points of D_v for all $v \notin \ell(x)$; therefore for $w \in W$, $\ell(w) \subseteq \ell(x) = \ell(b)$. Since $b \in B$, it must be in ∂D_z for some $z \in \ell(x)$. So there is a $w \in W$ such that $w \notin D_z$, thus $\ell(w) \subseteq \ell(b) \setminus \{z\}$, so the size of ℓ is not constant, a contradiction. $\square$

Theorem 17 has an interesting application to voting theory. In *approval voting*, each voter specifies which options she would consider acceptable, without ranking the options. Following [2], the set of all options available to voters is called a (*political*) *spectrum*; it often has a natural topology given by notions of “closeness” or “similarity” of political preferences. For instance, the political spectrum is often modeled as $\mathbb{R}$, a line with conservative positions to the right and liberal positions to the left. However, in elections over multiple issues, the spectrum might be best modeled as a subset of $\mathbb{R}^n$. Political spectra have been modeled also by a circle; often this arises by bending the linear political spectrum so that the extreme left-wing and right-wing positions are considered close; e.g., see [8].

For each voter, the set of options that a voter approves is called her *approval set*. We assume that approval sets are closed subsets of the spectrum, and we call the set of all voters together with their approval sets a *society*.

We call a society with a circular political spectrum *super-agreeable* if for each pair of voters i, j , one of the paths between i, j is covered by their approval sets. We remark that in many cases it is natural to assume that a voter’s approval set is connected. In this situation, a super-agreeable society is simply one in which every pair of voters can find common ground, i.e., an option which they will both approve. This agrees with the definition of super-agreeable for a linear society, as in [2].

Then Theorem 17 then has the following corollary:

Corollary 18. *In a super-agreeable society with a circular political spectrum, there is an option that will be approved by a strict majority of the voters.*

The value of this result is that it gives a sufficient condition for the existence of a strict majority using approval voting when the political spectrum is circular. We do not assume the approval sets have to be connected. Results for connected approval sets and weaker intersection hypotheses may be found in [2], who consider linear political spectra, and Hardin [5], who extends those results to circular political spectra.

REFERENCES

- [1] A. G. Aksoy and M. A. Khamsi. A selection theorem in metric trees. *Proc. Amer. Math. Soc.*, 134(10):2957–2966 (electronic), 2006.
- [2] Deborah E. Berg, Seguei Norine, Francis Edward Su, Robin Thomas, and Paul Wollan. Voting in agreeable societies. *To appear, American Mathematical Monthly*, 2010.
- [3] Eli Berger. KKM—a topological approach for trees. *Combinatorica*, 25(1):1–18, 2005.
- [4] James Dugundji and Andrzej Granas. *Fixed Point Theory*. Polish Scientific Publishers, Warszawa, 1982.
- [5] Christopher S. Hardin. Agreement in circular societies. *to appear, Amer. Math. Monthly*.
- [6] Mohamed A. Khamsi. KKM and Ky Fan theorems in hyperconvex metric spaces. *J. Math. Anal. Appl.*, 204(1):298–306, 1996.

- [7] B. Knaster, C. Kuratowski, and S. Mazurkiewicz. Ein beweis des fixpunktsatzes für n -dimensionale simplexe. *Fund. Math.*, 14:132–137, 1929.
- [8] Sven Gunnar Simonsen. *Nationalism and the Russian political spectrum: Locating and evaluating the extremes*. Studies on Foreign Policy Issues, 4:2000, Ministry of Foreign Affairs, Norway, 2000. Online at <http://odin.dep.no/ud/english/doc/reports/032001-220004/dok-bn.html>.
- [9] Mark Yoseloff. Topological proofs of some combinatorial theorems. *Journal of Combinatorial Theory (A)*, 17:95–111, 1974.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SAN DIEGO, LA JOLLA, CA 92093-0112

E-mail address: `aniederm@math.ucsd.edu`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CA 94720

E-mail address: `drizzolo@math.berkeley.edu`

DEPARTMENT OF MATHEMATICS, HARVEY MUDD COLLEGE, CLAREMONT, CA 91711

E-mail address: `su@math.hmc.edu`